

UNIVERSITA' CATTOLICA DEL SACRO CUORE
- Milano -

**QUADERNI DELL'ISTITUTO DI
ECONOMIA DELL'IMPRESA E DEL LAVORO**

The Indirect Effect of Fine Particulate Matter on Health through
Individuals' Life-style

Cinzia Di Novi

n. 62 – marzo 2011



QUADERNI DELL'ISTITUTO DI
ECONOMIA DELL'IMPRESA E DEL LAVORO

Istituto di Economia dell'Impresa e del Lavoro
Facoltà di Economia
Università Cattolica del Sacro Cuore
Largo Gemelli, 1 - 20123 Milano

* I quaderni sono disponibili on-line all'indirizzo dell'Istituto
<http://istituti.unicatt.it/ieil>

* Esemplare fuori commercio per il deposito legale agli effetti della
Legge n. 106 del 15 aprile 2004.

* La Redazione ottempera agli obblighi previsti dalla Legge n. 106
del 15.04.2006, Decreto del Presidente della Repubblica del
03.05.2006 n. 252 pubblicato nella G.U. del 18.08.2006 n. 191.

The Indirect Effect of Fine Particulate Matter on Health through Individuals' Life-style*

Cinzia Di Novi

Dipartimento di Politiche Pubbliche e Scelte Collettive, Università del Piemonte Orientale

Istituto di Economia dell'Impresa e del Lavoro, Università Cattolica del Sacro Cuore

E-mail: cinzia.dinovi@sp.unipmn.it

Abstract: Limited literature has been published on the association between environmental health indicators, life-style habits and ambient air pollution. We have examined the association of asthma prevalence and the amount of health investment with daily mean concentrations of particulate matter (PM) with a mass median aerodynamic diameter less than 2.5 μm ($\text{PM}_{2.5}$) in 16 metropolitan areas in U.S. using the Behavioral Risk Factor Surveillance System (2001) data in conjunction with the Air Quality System data collected by the Environmental Protection Agency. A multivariate probit approach has been used to estimate recursive systems of equations for environmental health outcome and life-styles. A piecewise linear relationship has been postulated to describe the association between health outcome, health investment and pollution. We have assumed one change point at AQI value of 100 which corresponds to the US national air quality standard. The most interesting result concerns the influence of pollution on health-improving life-style choices: below a specified threshold concentration (AQI=100) a positive linear association exists between exposure to $\text{PM}_{2.5}$ and health investments; above the threshold the association becomes negative. Hence, only if ambient pollution is in the 'satisfactory range' (AQI level at or below 100), individuals will have incentive to invest in health.

JEL-Classification: I12, C31, D13, D81, Q25

Keywords: health production, multivariate probit, spline, life-style, fine particulate, asthma.

* Earlier drafts of this study were presented at the 15th Italian Health Economics Association Annual Conference, Collegio Carlo Alberto, Moncalieri, Italy, the 20th Annual Conference of the Italian Public Economics Society (SIEP), University of Pavia, Italy. The author wishes to thank the participants of these conferences, in particular Silvia Balia, Vincenzo Carrieri, Massimiliano Piacenza, Silvana Robone, Gilberto Turati. The usual disclaimer applies.

1 Introduction

Under the 1970 Clean Air Act and subsequent amendments, introduced by the Environmental Protection Agency to limit the amount of air pollution, ambient air quality in the United States has improved dramatically. However, despite regulatory effort, fine particulate continues to be a matter for concern despite its falling level. The situation has been further aggravated by the fact that protection of public health is constrained by the inability of scientists to establish a safe level of $PM_{2.5}$ below which it poses little or no risks for human health. In fact, fine particulate even at much lower concentrations (below current US regulatory levels), has been associated with increased rates of mortality and morbidity in several cities in the United States, (in Europe and other developed countries, too) (Dume et al. 1998; Daniels et al., 2000; Bolin and Lindgren, 2002; Brunekreef and Holgate, 2002). Furthermore, the effect of particulate on health may be complex, as it may vary from one individual to another: scientists have to consider that individuals and groups are not equally vulnerable to air pollution health effects. Susceptibility factors could be strictly linked to fixed individual characteristics such as genetics, gender, age and race or to variable individual characteristics caused by the realities of life (Di Novi, 2010). Low socioeconomic classes, for instance, tend to be more susceptible to the adverse effects of air pollution because of other factors related to their life-styles: they are more likely to be uninformed over environmental health issues, to have an unhealthy diet, to smoke and drink alcohol, and in general to lead less healthy lives, with associated effects on their health (Grassman, 1996; Sexton, 1997). Hence, analysts must calculate changes in health outcomes by taking into account that the effect of pollution could easily be correlated with other factors that may be just as influential (Schwartz and Weiss, 1994 a, b; Brook et al., 2004)¹.

While, on the one hand, epidemiological studies have shown that pollution acts synergistically with tobacco smoking, alcohol consumption and unhealthy diet to induce respiratory illness such as asthma, lung cancer and cardiovascular diseases (Valavanidis et al., 2009) on the other hand there is little information on the extent to which quality of the environment may influence choices of life-style. This is an issue that has, in our view, received too little attention (see Cropper, 1981; Erbsland et al. 1994; Di Novi, 2010). An important contribution in this area was Cropper (1981), who explored the consequences of introducing pollution variables into the health production function. She considered changes in environmental conditions to influence the amount of health investments through the rate at which an individual's stock of health depreciates: Cropper assumes that when pollution increases, it becomes more costly to reduce the probability of a health shock. Individuals feel less healthy because they perceive the health depreciation rate to be higher. Hence, they may choose to invest less in their health and maintain lower health stock because of the higher net investment costs. In this sense, a higher pollution concentration may have two effects on health: a direct effect which consists of an increase of the health depreciation rate and an indirect effect, described by Cropper (1981), by which individuals will invest less in health and display a higher probability of suffering from bad health.

The purpose of this study is to examine the influence that the quality of the environment,

¹After age, height, body mass index, race, sex, cigarette smoking, and employment status were controlled for, Schwartz and Weiss (1994a, b) find that a diet rich in fish and vitamin C could have a protective effect on lung function; while cigarette smoking and heavy alcohol consumption interact with air pollution to increase mortality risks from cardiovascular diseases (Brook et al., 2004).

captured by the $PM_{2.5}$ level, may have on health investment decisions. The paper divides into two parts. The first part provides theoretical framework built on the basic concepts and ideas of the demand for health by Grossman (1972) and the subsequent contribution by Cropper (1981). In its second part the paper provides empirical support to the theoretical assumptions.

In the empirical part of the paper, in order to introduce a measure of health stock, a dichotomous measure of asthma prevalence has been used. We choose asthma since it simultaneously represents a health outcome and an "environmental health indicator" (see WHO, 1999). Since we have included life-style variables as regressors in the health equations, a problem of simultaneity may arise. Hence, we try to correct the potential endogeneity of the behavioral variables by using a recursive multivariate probit model which is available in the literature although not so frequently used (Blaylock and Blisard, 1992; Contoyannis and Jones, 2004; Balia and Jones, 2008; Di Novi, 2010). A piecewise linear function has been employed to describe the relationship between health, health investments and pollution. We assume one change point at AQI value of 100 which corresponds to the U.S. national air quality standard.

The model is estimated using data from the 2001 Behavioral Risk Factor Surveillance System (BRFSS). BRFSS does not measure environmental quality but it can be used in conjunction with the 2001 Environmental Protection Agency's (EPA) Air Quality System (AQS) database. We merged data from the AQS with BRFSS data using the metropolitan area information. The EPA's Air Quality System (AQS) database contains measurement of six criteria pollutants: ozone (O_3), sulphur dioxide (SO_2), carbon monoxide (CO), and particulate matter ($PM_{2.5}$, PM_{10}). Because our study focused on $PM_{2.5}$ we used the daily AQI which reported daily air quality based on the concentration levels of $PM_{2.5}$.

The most interesting-and possibly surprising- result is the effect that pollution appears to have on health-improving life-style choices. This result partly contradicts what one should expect from Cropper's model, where pollution makes the investments in health more costly. In order to rationalize the empirical result obtained, one should refer to the relationship between pollution and the investments in health as an inverse-V-shaped emission-health investments relationship with a threshold pollution point: only if air pollution is concentrated above this point individuals will no longer have incentives to invest in health-improving activities. This result may have an important policy implication: an intervention that reduces air pollution below the threshold pollution level, may have not only a direct effect on individuals' health status, but also an indirect health effect through a healthier life-style which seems to be one of the driving factors for good health.

The rest of the paper is organized as follows: section 2 introduces a model of health production. Section 3 describes the data and the variables for the analysis. Section 4 presents the empirical approach and the econometric results. Section 5 concludes with a discussion. The definition of the variables, descriptive statistics and tables with estimation coefficients are in Appendix.

2 A Model of Health Production

We assume that each individual is endowed with a stock of health capital H_t that evolves according to:

$$\Delta H = H_{t+1} - H_t = f(\Lambda(E), t) - \delta_t H_t - \mathcal{G}_t \quad (1)$$

where $\delta_t \in (0, 1)$ is the natural rate at which health deteriorates. \mathcal{G}_t is a random shock. We assume that the shock could be any injury which causes a reduction in the current state of health. Moreover, we assume that \mathcal{G}_t can take a value of zero when the shock does not occur and a positive value $\mathcal{G}_t > 0$ when it does occur. The transition probability of having a shock next period is assumed to be inversely related to the stock of health. Then, the size of health is important since it affects the probability for an individual of enjoying good or bad health. Individuals can affect the probability of bad or good health next period by "investing" or "disinvesting" in health. The investments/disinvestments in health are captured by a household production function $f(\Lambda(E), t)$. Where Λ indicates the individuals behavior. We distinguish between healthy and unhealthy behavior. A proxy for healthy behavior consists, for instance, in a healthy diet (fruits and vegetables consumption etc.) or in sport activities practice, while a proxy for unhealthy behavior includes consumption of hazardous goods like alcohol consumption or cigarettes smoking. E is the exogenous education level that is assumed to affect the productivity of producing health. Schooling helps people choose healthier life-styles by improving their knowledge of the relationship between health behaviors and health outcomes (Berger and Leigh, 1989; Kenkel, 1991). A more educated person may have more knowledge about the harmful effects of cigarette smoking, pollution exposition, alcohol consumption or about what constitutes an appropriate, healthy diet. Furthermore, schooling increases information about the importance of having regular exams or screening tests to prevent an illness or at least to minimize disease. $f(\Lambda(E), t)$ can increase or fall in individual behavior Λ . In particular $f(\Lambda(E), t)$ is increasing in a healthy behavior and decreases if individuals disinvest in their health by consuming, for instance, hazardous goods. It follows that while a healthy life-style increases the stock of health capital, actions detrimental to health such as cigarette smoking and excessive alcohol consumption lower the stock of health capital.

In order to introduce the impacts of the environment, our analysis takes changes in environmental conditions to influence the rate at which an individual's stock of health depreciates:

$$\delta_t = \delta_0 (1 + \tilde{\delta}) \Psi_t^\phi \quad (2)$$

Following Grossman (1972) and subsequent contribution by Cropper (1981) we assume that health depreciates over time at an increasing rate with age ($\tilde{\delta}$) and with the ambient air pollution to which an individual is exposed (Ψ). Pollution enters directly the rate of decay and physically alters the state of a person's health; its effect is measured by ϕ .

As in Cropper's (1981) model, we assume that the individual behavior is influenced by environmental pollution. We assume that there is an optimal pollution level Ψ^* to maximize health investments and healthy behaviors. An increasing level of pollution encourages health investments if it does not exceed a certain threshold. But if pollution level exceed the optimal

threshold, a decrease of ambient air quality may lead individuals to invest less in health. Individuals may have no incentives to invest in health since they feel less healthy because they perceive δ to be higher. Hence, they may choose to invest less in their health and maintain lower health stock because of the higher net investment costs. In this sense, a higher pollution concentration above the optimal threshold may have two effects on health: a direct effect which consists in an increase of δ and an indirect effect, described by Cropper (1981), by which individuals will invest less in health and display a higher probability of suffering from health shocks. In Section 4, we will provide empirical support to this theoretical assumption and we will test if there exist an optimal threshold in the relationship between pollution and life-style variables.

3 Data and Variables

Data pertaining to health status and health-related characteristics were drawn from the 2001 Behavioral Risk Factor Surveillance System Survey². The BRFSS is an ongoing state-based, landline cross-sectional telephone survey used to collect information on health risk behaviors, preventive health practices, access to and use of health care services primarily related to chronic conditions among U.S. adults aged 18 years or older. BRFSS does not directly measure environmental quality, information at the metropolitan area-level is available from the 2001 EPA's Air Quality System (AQS) database and can be used in conjunction with BRFSS. Thus, we have merged the two data sets by metropolitan area³.

The term metropolitan area is a geographic subdivisions formally defined by the U. S. Office of Management and Budget (OMB), for use by Federal statistical agencies in collecting, tabulating, and publishing Federal statistics. Following the OMB definition, the metropolitan area is "*an area containing a recognized population nucleus and adjacent communities that have a high degree of integration with that nucleus*" (OMB, 2000). The term metropolitan area refers collectively to metropolitan statistical area (MSA), primary metropolitan statistical area (PMSA), or New England county metropolitan area (NECMA):

1. a MSA consists of one or more counties with a high degree of social and economic integration, the presence either of a city with 50,000 or more inhabitants, or an urbanized area (i.e. a contiguous area of relatively high population density), and a total population of at least 100,000;
2. a PMSA consists of one county or group of counties that qualify as an MSA and have a total population of 1 million or more;
3. because of the greater importance of towns over counties, for the New England states, metropolitan areas are defined by a collection of towns and cities instead of by counties with a presence a total population of at least 75,000.

We have excluded respondents with missing air pollution information as well as those with an unknown metropolitan area of residence. Finally, we have dropped all those living outside the continental USA. After correcting for missing values, the final sample contains

²Centers for Disease Control and Prevention (CDC). Behavioral Risk Factor Surveillance System Survey Data. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention (2001).

³Because the environmental data from the AQS were collected during 2001 and reported at the metropolitan area-level, they should only be used in conjunction with 2001 BRFSS data that have been re-weighted for metropolitan area-level analysis.

6,760 individuals distributed in 16 metropolitan areas: Chicago, IL; Cincinnati, OH-KY-IN; Cleveland-Lorain-Elyria, OH; Columbus, OH; Des Moines, IA; Huntington-Ashland, WV-KY-OH; Las Vegas, NV-AZ; Louisville, KY-IN; Memphis, TN-AR-MS; Milwaukee-Waukesha, WI; Minneapolis-St. Paul; MN-WI; Nashville, TN; Omaha, NE-IA; Phoenix-Mesa, AZ; St. Louis, MO-IL; Tucson, AZ. Here, the term MSA refers to PMSAs.

Each respondent in the BRFSS was assigned an average annual $PM_{2.5}$ exposure based on his or her metropolitan area of residence⁴. Exposures are measurements taken at a network of national, state and local air monitoring stations which report the concentrations for $PM_{2.5}$. As $PM_{2.5}$ is measured in each metropolitan area at multiple sites, an annual mean has been computed and has been used in the analysis.

The EPA's Air Quality System database contains $PM_{2.5}$ AQI measurements. Formerly known as the Pollutant Standard Index (PSI)⁵, the nationally uniform AQI is a single number used by state and local agencies for reporting the air quality with respect to its effects on the human health. To make the AQI as easy to understand as possible, EPA has divided the AQI into six categories that correspond to different level of health concern with a scale that runs from 0 to 500 associated with a color scheme that runs from green to maroon:

1. 0-50 (green)- good, i.e. air quality poses little or no risks;
2. 51-100 (yellow)- moderate, i.e. air quality is acceptable but may be some concern for very small number of people;
3. 101-150 (orange)- unhealthy for sensitive groups, i.e. members of sensitive groups (i.e. children, older adults, people with respiratory disease or heart disease) may experience health effects. The general public is not likely to be affected;
4. 151-200 (red)- unhealthy, i.e. everyone may begin to experience health effects;
5. 201-300 (purple)- very unhealthy, i.e. everyone may experience more serious health effects;
6. 300-500 (maroon)- hazardous, i.e. the entire population is more likely to be affected and everyone should avoid all outdoor exertion.

In 1976 the U.S. EPA established the National Ambient Air Quality Standards (NAAQS) of 100: only if the AQI level is at or below 100 ambient air quality can be considered in the satisfactory range.

3.1 Environmental Health Indicator and Life-Style Variables

As "environmental health indicator" (WHO, 1999) a dichotomous measure of asthma prevalence has been used. Asthma is one of the most common chronic illness in the world and represents one of the most important cause of morbidity, economic cost and mortality in developed and developing countries alike: approximately 300 million people currently suffer from asthma worldwide. In North America about 1 person in 10 has been diagnosed with asthma. The asthma prevalence rates in the United States are approximately 10.9%, representing

⁴An important limitation of our analysis concerns the particulate matter data which provide estimates for ambient air particulate matter at the metropolitan area level for a one-year period and not at an individual exposure level to the daily concentrations level.

⁵Based on available evidence that fine particles were particularly damaging, in 1999 the PSI was revised by US Environmental Protection Agency and replaced by the Air Quality Index (AQI) to incorporate new $PM_{2.5}$.

35.5 million individuals (Masoli et al. 2004; Braman, 2006; Bateman et al., 2008;). Asthma has been associated with worse health-related quality of life among all adult (Ford et al. 2003).

Although understanding of many aspects of asthma has improved over the past decades, the fundamental causes of the tendency to develop asthma remains largely unknown (Braman, 2006). The development of asthma seems to be related to certain asthma genetic factors and to individual exposures to certain life-style (in particular to smoking habits and obesity) and environmental factors (allergen exposure, tobacco smoking, outdoor air pollution) (Basagaca, et al., 2004; GINA, 2006). Concerning outdoor pollution, there is evidence that long term exposure to substances found in the air, such as particles, may interact with genetic factors to develop and to determine the subsequent maintenance of asthma (Bascom et al.,1996; Katsouyanni et al. 1997; Brunekreef and Holgate, 2002). In order to capture the effect of pollution on asthma,we have introduced a self-reported measure of asthma prevalence. To be classified as having asthma, the respondent must have answered "yes" to the following two fairly precise questions: "*Have you ever been told by a doctor, nurse, or other health professional that you have asthma?*"and "*Do you still have asthma?*".

The life-style variables⁶ indicate whether the individual is a smoker, consumes heavy alcohol, eats fruits and vegetables at least once per day, meets recommendations for physical activity⁷, is obese, suffers from mental stress and depression, has had flu shot vaccination during the year of the interview⁸. We use BMI (Body Mass Index) to compute an indicator of obesity⁹ according to medical and WHO guidelines. Table 1 contains the full description of the life-style variables¹⁰.

TABLE 2a) shows a simple descriptive analysis, which presents sample means and standard deviations for the variables used in the models. It is worth noting that the sample (40.7 % men) consists of individuals whose behaviors are mostly healthy with the exception of the variables that assess physical activity: only 25 per cent of individuals are current smokers, 5 per cent of individuals drink heavily, and 36 per cent of them suffer from stress; while 96 per cent of them follow a healthy diet and 22 per cent devote some time to physical activity.

3.2 Other Characteristics

The other independent variables in the model can be grouped into categories which are

⁶The endogenous behavioral variables employed are those which cover as much as possible the life-style categories used by Belloc and Breslow's (1972) epidemiological study based on the Alameda County survey carried out in California in 1965.

⁷Based on the "Physical Activity Guidelines for Americans" (2008) exercise recommendations for healthy adults and older adults are 300 minutes a week of moderate-intensity, or 150 minutes a week of vigorous-intensity aerobic physical activity, or an equivalent combination of moderate- and vigorous-intensity activity (Physical Activity Guidelines for Americans, 2008).

⁸ Influenza is an infectious disease with possible severe health consequences for the elderly (over 65) and non-elderly in poor health (Mullahy, 1999). The vaccination of adults and children with asthma or reactive airway disease is recommended every year by the Advisory Committee on Immunization Practices (ACIP) since they are at increased risk of complications from influenza infection (Advisory Committee on Immunization Practices, 2009).

⁹Obesity is considered a risk factor for several diseases. It is often associated with aspects of an individual's life-style such as insufficient exercise and inappropriate diet or nutrition. Those who are obese are expected to have poorer health

¹⁰The original variables are mostly discrete qualitative indexes but with more than two modalities; clearly it would be better to use this information, but the computational complexities would have increase considerably with uncertain benefits for the point of view of the estimation quality.

listed, together with the life-style variables, in TABLE 1. We consider the following categories: physical characteristics such as age, sex, race, prior health in order to capture health status at the beginning of the observation period¹¹; a measure of leisure time physical activity (as a predictor of healthy behaviors); household composition; air pollution; socioeconomic variables such as health coverage (including HMO plan¹²), education, marital status, employment status. Despite many research findings have suggested a significant associations between income inequality and health, we do not include income among socioeconomic variables because of the unreliability of this information¹³.

Concerning leisure time physical activity, the following item was used to create a binary outcome: "During the past month, did you participate in any physical activities such as running, calisthenics, golf, gardening, or walking for exercise". The response was "yes" or "no". Individuals answering "no" to this question were categorized as inactive, and those answering "yes" were categorized as "active". Leisure time physical activity has been shown to be a strong predictor of regular moderate physical activity or physical fitness necessary to improve health (Sandvik et al., 1993; Dishman and Sallis, 1994). In our sample, of those who reported leisure time physical activity (74.8 %), 28.22% meet recommendations in physical activity, while among those who initially reported no leisure-time physical activity (25.2%) only 3.99% exercise to meet health-related recommendations (Physical Activity Guidelines for Americans, 2008).

Recent studies have found that the absence of leisure time physical activity is a good measure for the tendency to sedentary behavior that is considered an important reason for obesity and for poor dietary habits: people who are active generally have healthier dietary habits than their sedentary counterparts (Sherwood and Jeffery, 2000). Moreover, sedentary behavior is strictly linked to smoking behavior and to the use of other substances as well, such as alcohol or high-fat foods (King and Brassington, 1997; Marcus et al., 1999).

The main variable of interest in the model is fine particulate. Particulate matter (PM) is an air pollutant consisting of dust, dirt, soot, smoke, and liquid droplets suspended in the air. Particles with aerodynamic diameters between 2.5 and 10 micrometers consist mainly of crustal particles mechanically generated from agriculture, mining, construction, road traffic, and related sources, as well as particles of biological origin. While particles with aerodynamic diameters 2.5 consist mainly of combustion particles from motor vehicles and the burning of coal, fuel oil, and wood, but also contain some crustal particles from finely pulverized road dust and soils. PM has been associated with a wide range of related human adverse health effects, including an

¹¹We have included a measure of prior physical health with a time frame of 30 days, refers to a measure of self-reported physical health: respondents were asked if their physical health was not good in the thirty days before the interview.

¹²A health maintenance organization (HMO) is a type of managed care plan that provides health coverage in the United States to its members through a network of doctors, hospitals, and health care providers. HMOs are popular alternatives to traditional health care plans offered by insurance companies because they can cover a wide variety of services, usually at a significantly lower cost.

¹³The BRFSS does not collect a continuous measure for income. The only measure available in the BRFSS is household income which is provided in the form of eight intervals, with the highest category listed as "\$75,000 or more". Respondents are assigned a value corresponding to the midpoint of their income interval. In general, missing data is not a large problem with the BRFSS questions: response is around 95% or higher for all questions with the exception of baseline household income which suffer a high rate of item non-response (around 20%). Because of the high rate of non-response, we have not included income given the potential for sample selection bias and the reduction in sample size induced by using only those individuals for whom we have this information.

increased incidence of chronic bronchitis and acute respiratory illnesses, exacerbation of asthma, impairment of lung functions and premature mortality (Dume et al. 1998; Bolin and Lindgren, 2002; Brunekreef and Holgate, 2002). The size of particles is directly linked to their potential for causing health problems (EPA, 2003). In particular smaller or fine particles ($PM_{2.5}$) seem to pose the greatest problems since they may penetrate more deeply into the lung and may reach the alveolar region.

4 Empirical Approach

4.1 Multivariate Estimation

From a methodological point of view, it should be noted that asthma equation is a structural equation since the health behavior inputs may be endogenous. Efficient and consistent estimation of the parameters in the health equation requires a model that takes account of the nature of the variables used. The potential simultaneity, which can arise with the inclusion of life-style variables as regressors, can be corrected by using a recursive multivariate probit model¹⁴ (Contoyannis and Jones, 2004, Balia and Jones, 2008; Di Novi, 2009).

We identify two classes of dependent variables: the individual health behaviors and the health outcome. The seven equations for the health behavior variables are modeled as reduced-form equations. The asthma equation is the structural equation with the health behavior variables as explanatory factors. Hence, we construct and estimate a system of eight equations ($m = 8$) with seven reduced-form and one structural equation.

Thus:

$$\begin{aligned}
 y_{li}^* &= b_1' x_{li} + e_{li} = d_2 y_{2i} + d_3 y_{3i} + d_4 y_{4i} + d_5 y_{5i} + d_6 y_{6i} + \\
 &+ d_7 y_{7i} + d_8 y_{8i} + a' z_i + f(PM_{2.5} AQI) + e_{li} \\
 &\times \\
 &\times \\
 &\times \\
 y_{li}^* &= b_l' x_{li} + f(PM_{2.5} AQI) + e_{li}
 \end{aligned} \tag{3}$$

where x_{li} (with $l = 2, \dots, 8$) and z_i are vectors of exogenous variables, β_l and α are parameter vectors, δ_l is a scalar parameter and $\beta_l' = (\delta_l, \alpha')$. $f(PM_{2.5} AQI)$ is some function of $PM_{2.5} AQI$. Two forms of $f(PM_{2.5} AQI)$ has been used in this article:

- linear function $\eta PM_{2.5} AQI_i$; and
- piecewise linear function with one change point:

¹⁴The multivariate probit model with endogenous dummies belongs to the general class of simultaneous equation models (see Maddala, 1983).

$$f(PM_{2.5} AQI) = \begin{cases} \eta_1 PM_{2.5} AQI_i & \text{if } PM_{2.5} AQI \leq \theta \\ \eta_1 PM_{2.5} AQI_i + \eta_2 (PM_{2.5} AQI - \theta)_i & \text{if } PM_{2.5} AQI > \theta \end{cases}$$

where η_n (with $n=1,2$) are scalar parameters, and θ is the change point fixed at 100 which corresponds to the national air quality standard i.e. the $PM_{2.5}$ AQI value at which the slope of the piecewise linear function is allowed to change (see Robert, 2004).

$\varepsilon_{1i}, \dots, \varepsilon_{li}$ are the error terms distributed as multivariate normal, each with a mean zero and a variance covariance matrix Σ . Σ has values of 1 on the leading diagonal and correlations $\rho_{jk} = \rho_{kj}$ on off-diagonal elements (where ρ_{jk} is the covariance between the error terms of equation j and k).

In the above setting, the exogeneity condition is stated in terms of the correlation coefficient, which can be interpreted as the correlation between the unobservable explanatory variables of the different equations. All the equations in (3) can be estimated separately as single probit model only in the case of independent error terms i.e. the coefficient ρ_{jk} is not significantly different of zero.

The parameters of the equations are not identified if z_i includes all the variables in x_{li} . Estimation requires some considerations for the identification of the model parameters. Maddala (1983) proposes that at least one of the reduced-form exogenous variables (x_{li}) is not included in the structural equations as explanatory variables. Following Maddala's approach we impose exclusion restrictions. For the reduced form, we use marital status¹⁵ assuming that it has only an indirect effect on health through the life-style variables. Moreover, in order to address the identification problem, we include in the reduced form equation the measure of leisure time physical activity. In addition, we exclude from the asthma equation the variables that indicate the number of adults and children living in the household which are considered to influence to a certain extent individual's preferences and decisions about health¹⁶.

The reference individual in the model is female, married and employed. She is aged eighteen years old or more and she has attended elementary school. The estimation of a multivariate probit is carried out using the STATA 10 software which applies the method of Simulated Maximum Likelihood estimation (see Cappellari and Jenkins, 2003). STATA provides the statistic $z = \hat{\rho}/S_{\hat{\rho}}$ to test the hypothesis $H_0: \rho = 0$. If the error terms are independent, the Maximum Simulated Likelihood estimation is equivalent to the separate Maximum Likelihood probit estimation.

¹⁵To balance statistical fit of the model we use the Bayesian information criterion proposed by Schwarz (1978). This criterion suggests the exclusion of the variables that describe marital status from the health outcomes and the SAH equation. Kenkel (1991), Contoyannis and Jones (2004), Balia and Jones (2008) exclude marital status from the health and the death equation claiming that marital status influence only indirectly the probability of good or bad health and the probability of death, through the life-style habits: smoke, alcohol, diet etc.

¹⁶Family structure and composition may affect health behaviours. Takeda et al. (2004), for instance, find that an increasing number of women in households is associated with a strong presence of protective health behaviours (less smoking, less heavy drinking), but also with more sedentary behaviour, while the presence of men in household are associated with a higher probability of heavier smoking. Moreover Bakhshi et al. (2008)'s paper shows that there exists an association between the number of young children and obesity among men and women aged 20-75 years.

4.2 Estimation Results

The first column of TABLE 3a) and TABLE 3b) shows coefficients for the structural asthma equation estimated in the full recursive model, using the multivariate probit specification.

Starting from life-style variables, we can observe that the results support the theoretical model: in the asthma equation smoking behavior, alcohol consumption, being stressed and being obese have the expected significant positive influence on the probability of suffering from asthma. Physical activity and diet variables do not contribute to explain the probability of reporting asthma. Concerning the preventive medical behavior, flu vaccination shows a positive and statistically significant coefficient in the asthma equation. One of the possible reasons for the ambiguous sign is that persons getting flu shots are more likely to have physicians and this may in turn lead to greater detection of the disease than in those who do not have regular medical care access.

Medical literature clearly indicates that asthma tends to be a disease of young people: the prevalence of asthma is highest in people younger than age 18 and tends to decrease with increasing age. Our model too, predicts that the probability of suffering from asthma decreases with age. Also gender affects the probability of suffering from asthma: according to the medical literature, in our sample asthma seems to be more prevalent in women (Dicpinigaitis and Rauf, 1998; De Marco et al. 2000; Eagan, et al., 2002).

Concerning the effect of pollution, first a linear function and then a piecewise linear function were fitted to fine particulate using the procedure *mkspline* function¹⁷ of STATA 10. While the relationship between $PM_{2.5}$ and asthma seems to be adequately described by a linear model¹⁸, with a positive statistically significant effect on asthma prevalence, the effect of fine particulate on individuals' health-improving activities seems to be better represented in an inverse-V-shaped emission-health investments relationship with a threshold pollution point (see Figure 1¹⁹). The changing point at which the slope of the piecewise linear relationship changes is the AQI value of 100²⁰. From our empirical results, it arises that when fine particulate is concentrated above national air quality standard, people start to disinvest in health. Above the

¹⁷ The *mkspline* command creates the linear spline variables $pm2.5_aqi < 100$ and $pm2.5_aqi > 100$. The coefficients for these variables give the slope in each of the regions before and after the knot, which we placed at $PM_{2.5}$ AQI value of 100. When the marginal option is specified, the new variables are constructed so that, when used in estimation, the coefficients represent the change in the slope from the preceding interval. However, the default is to construct the variables so that the coefficients will measure the slopes for the intervals rather than the difference in the slopes. So, we do not use marginal option (see Gould, 1993).

¹⁸ When the relationship between particulate matter concentration and health indicators have been evaluated, most studies have reported no evidence of a clear threshold concentration below which the harmful effect of fine particulate is less likely (Samet et al. 1995; Daniels et al., 2002; Dominici et al., 2002). According to these studies we can conclude that the relationship could be reasonably considered linear.

¹⁹ Figure 1 shows, in raw form, the relationship between health investments and pollution.

²⁰ An AQI value of 100 generally corresponds to the national air quality standard for the pollutant, which is the level that the EPA has set to protect public health. AQI values below 100 are generally thought of as satisfactory. When AQI values are above 100, air quality is considered to be unhealthy at first for members of sensitive populations (i.e. children, elderly and people who suffer from respiratory or cardiovascular diseases) then for everyone as AQI values go above 150. Many US larger cities have severe air pollution problems, and the AQI in these areas may exceed 100 even though AQI values higher than 200 are infrequent, and AQI values above 300 are extremely rare. For the metropolitan areas under consideration in this study $PM_{2.5}$ AQI displays values between 62 and 161 (with the higher value experienced by Milwaukee-Waukesha, WI).

"PM_{2.5} AQI optimal point" if air pollution level increases along the downward-sloping portion of the curve, individuals will have no incentives to invest in health. While a reduction in fine particulate along the downward-sloping portion of the curve may lead to an increase in the health-improving life-style choices. Thus, it arises that, concerning the effect of pollution on health investment decisions, the most important number in the AQI scale is 100²¹ (the standard established under the Clean Air Act): only if the PM_{2.5} AQI level is at or below 100, fine particulate will be in the satisfactory range and people will have an incentive to invest in health. According to our results while a higher concentration of PM_{2.5} when fine particulate is in the satisfactory range would have a positive influence on healthy habits (in particular, a negative influence on smoking behavior, alcohol consumption, stress, and a positive effect on diet and flu vaccination) when PM_{2.5} AQI values go above 100 an increasing level of fine particulate seems to lead individuals to invest less in health-improving activities with a positive effect on the probability of smoking, consuming heavy drinks and suffering from stress; in addition, it decreases the probability of following a diet rich in fruits and vegetables and having preventive care²². Our model predicts that people increase physical exercise in response to higher PM_{2.5} level when pollution is outside the satisfactory range. However, this does not always seem to be a healthy choice, in particular when individuals exercise outdoors: when AQI values exceed 100, air quality is considered to be unhealthy, and is always accompanied by EPA advice to avoid harmful air pollution and especially those with medical conditions affecting the heart and lungs should consider limiting the type, duration and location of outdoor activity pursuing alternative physical activities, such as indoor activities. Activity outdoors at times when the air is polluted can harm the heart and lungs (Wen et al. 2009) and may lead to other serious health effects.

Concerning socioeconomic indicators, it arises that being outside the labor force (students, retired and homemaker) has a positive influence on developing asthma. This result can be explained by the fact that homemakers, students and retired people spend more time indoors (95% of their time against 90% of their employed counterpart - 2008 American Time Use Survey) where the quality of air is often poor since it may be affected by indoor pollution (i.e. environmental tobacco smoke and other indoor allergens) that even at relatively low levels, may play a significant role in the development of asthma (Jones, 1998; Ernst, 2002). On the other hand, people outside the labor force show a higher probability of following healthier behaviors, with respect to the employed and those who are involuntarily unemployed: they smoke less, they consume heavy drinks less frequently and suffer less often from stress and obesity than people who are in the labor force (even though they more often tend to settle down to a more sedentary life). The other socioeconomic indicators seem to have no influence on asthma prevalence. In the empirical literature, social class (based on occupation, education and

²¹ We carried out a sensitivity analysis re-running the model with different thresholds for PM_{2.5}. A one change piecewise linear model was fitted fixing the change point at AQI=150. The model was fitted again using linear splines with knots at AQI=100 and AQI=150. AIC information criterion was computed for each model. The model with the smallest AIC value was chosen as the "best" (Roberts, 2004). The model with one change point at AQI=100 had the smallest AIC value, so this was used as evidence to support the conclusion that the relation between health investments and PM_{2.5} was adequately described by a piecewise linear model with one change point at AQI=100. For the sake of brevity, the results of the sensitivity analysis are not included but they are available on request.

²² These results confirm the findings of Di Novi (2009).

income) has been suggested to relate to asthma: some studies have shown increased asthma hospital admissions and asthma severity in low social class groups (see, for instance, Walters et al., 1995; Watson et al., 1996; Earzen et al., 1997; Chen et al. 1999), but the association between socioeconomic deprivation and asthma prevalence is less clear and consistent evidence is still lacking. The tendency to develop asthma seems to be more related to certain asthma genetic factors (for instance, a family history of asthma may contribute to the development of asthma) and to individual exposure to certain life-style and environmental factors (Basagaca et al., 2004). On the other hand, socioeconomic status seems to affect health improving activities. There is a clear indication of the allocative effects of schooling, since schooling is related to life-style variables in a health promoting way: on the one hand, attending a college or being a college graduate has a negative impact on cigarettes and alcohol consumption, on stress, and on the risk of obesity. On the other hand, a higher degree of education is positively related to the probability of having the flu shot, meeting physical activity recommendations and following a healthy diet.

Our results do not offer significant evidence of racial disparity in asthma prevalence between White and Black Americans. The disparity in asthma prevalence and treatment between races has been studied at length. From the literature it arises that the disparity in asthma hospitalization is greater than the disparity in asthma prevalence, which suggests that once asthma is established, the black/white gap in asthma seems to be explained by other factors (such as differential access to medical care, differential access to housing, differential patterns of medical care use). Instead our results show a black/white gap in obesity: being black increases the likelihood of obesity. Obesity has reached near epidemic proportions in the United States (Burkhauser et al. 2009). Together with cigarette smoking, obesity is one of the leading causes of several chronic conditions and mortality in U.S. The prevalence of obesity is high among African Americans, particularly African American women (Stolley and Fitzgibbon, 1997).

As for the household composition variables, we can observe that the presence of children younger than eighteen years old is negatively correlated with alcohol consumption, but it has a negative influence on exercise and on preventive care and increases the probability of being obese. An increasing number of women in a household increases the probability of choosing a healthier diet and of reducing cigarette smoking, but has a positive impact on stress and a negative influence on exercise indicators.

Many empirical findings have documented a potential health benefit of marriage: married people (including those who cohabit) appear to be healthier and to have a longer life expectancy than the unmarried (Espinosa and Evans, 2008). Some of the most convincing evidence is consistent with the marriage protection hypothesis, which assumes that "... married individuals engage in low-risk activities, share resources and enjoy caring for each other..." (Hu, Wolfe, 2002). Our model is consistent with the previous findings: in fact, from the empirical results, it arises that marital status has a significant influence on the life-style variables. In particular, marriage seems to be positively correlated to healthy habits while being divorced, separated, never married, or an unmarried couple has a positive impact on smoking and on alcohol consumption and on bad habits in general.

As mentioned previously, we have estimated the eight equations together using the multivariate probit specification. The multivariate probit allows us to test for unobserved heterogeneity whose effect is captured by the correlation between the error terms from the single equation models. By estimating all eight equations at the same time and taking account of

correlation in the error terms for the eight equations we are able to control for the effect on unobserved factors.

TABLE IV shows the correlation matrices for the full recursive models. The null hypothesis of exogeneity is rejected in eleven cases in the linear model and in nine cases in the piecewise linear model²³. Starting from the asthma equation, the correlation parameters indicate whether and how unobservable factors jointly affect life-style decisions and health outcome. As we can note, there exists a negative statistically significant correlation between the disturbance of the asthma equation and the equation for alcohol consumption and the obese equation. The negative coefficient concerning alcohol and obese support weak evidence that individuals with poor health tend to have a healthier life-style.

From the correlation matrices it appears that unobservable factors, which affect the propensity to smoke are positively related to those that affect heavy drinks consumption and being stressed, and it is negatively related to the propensity of consuming fruit and vegetables. Alcohol consumption is positively related to stress and negatively related to a healthy diet. From these results, it emerges that the unobserved propensity for a unhealthy life-style seems to be explained by unobserved characteristics which determine whether an individual smokes and consumes alcohol.

5 Summary and Conclusions

Limited literature has been published on the association between environmental health indicators, health improving activities and ambient air pollution. We have examined the associations of asthma prevalence and the amount of health investment with daily mean concentrations of fine particulate ($PM_{2.5}$) in 16 metropolitan areas in U.S. using the Behavioral Risk Factor Surveillance System (2001) data in conjunction with the Environmental Protection Agency's Air Quality System data.

The theoretical framework is based on a Grossman (1972)-style health production model and subsequent contribution by Cropper (1981) set up in discrete time.

A multivariate probit approach has been used to estimate recursive systems of equations for environmental health outcome and life-styles. We have assumed that the relationship between investment in health and ambient air quality could be represented by an inversely V-shaped curve, which assumes the existence of an optimal pollution level at which health investments are maximized. Based on this assumption, a piecewise linear relationship has been employed to describe the association between health investment and pollution. We have assumed one change point at AQI of 100 which corresponds to the national air quality standard. The empirical results show that below the specified threshold concentration (AQI=100) a positive linear association exists between exposure to $PM_{2.5}$ and health improving activities: people may invest more in health to offset the higher decline rate. But, above the threshold the association becomes negative: if particulate concentration is above the AQI optimal level, an increase in $PM_{2.5}$ along the downward-sloping portion of the curve may lead individuals to invest less in health, since it becomes more costly to build up resistance against pollution.

²³The statistically significant correlation coefficients suggest that the null hypothesis of eight univariate probit models or the hypothesis of independence across the error terms of the eight latent equations, can be rejected, and the multivariate probit model is a better model for the observed data.

Because of the higher investment costs, individuals may have lower incentives to invest in health and they may choose to maintain lower health stocks. Hence, only if ambient pollution is in the 'satisfactory range' (AQI level at or below 100), individuals will have incentive to invest in health. In this sense, a pollution concentration above the national air quality standard may have two effects on health: a direct effect which consists in an increase of health deterioration rate (with negative consequences on health stock) and an indirect effect (the same described in Cropper, 1981), by which individuals will invest less in health and display a higher probability to suffer from health shocks.

In urban areas, diesel vehicular traffic, often one of the main contributors to pollution including highly damaging emissions of particulate matter which fall into the fine particulate range, is more dangerous for human health (in contrast to coarse particles). Hence, from a policy perspective, intervention that combines public education (for instance, by increasing the number of passengers per vehicle, reducing trips on poor air quality days, keeping vehicles well-maintained, purchasing and using low emission vehicles, using alternative fuels etc.) with other action that abates pollution in the transport sector, would be a key part of urban air quality management strategies in order to reduce fine particulate to below the "optimal threshold level"; such intervention may have not only a direct effect on individuals' health status, but also an indirect health effect through a healthier life-style which is, based on our theoretical and empirical results, one of the driving factors for good health.

This paper suffers from two major limitations: first, we have established a model of health production which is dynamic to show the theoretical relationship between individual life-styles and the rate of health deterioration due to air pollution. Evidently, cross-sectional design of this study involves the evaluation of the characteristics of the individuals and their health and health behaviors at the same point in time making it impossible to disentangle the temporal sequencing of individual behaviors and failing to take the dynamic nature of the health accumulation model into account. In order to sort out these temporal sequences, long-term studies are needed. However, data of this type are rarely available. Moreover, where panel data do exist, they often lack the details on environmental or personal behaviors- such as cigarette smoking, alcohol consumption, dietary habits, and physical inactivity. So, despite the limitations of the cross-sectional data, we have used BRFSS survey since it provides rather detailed information about health status, diseases, life-style, education and other individual characteristics and it can be combined with the environmental information available from other sources, such as the Environmental Protection Agency and state and local monitoring networks, to compare measures of environmental quality and chronic disease.

Second, in estimating the relationship between particulate and health, we have used ambient levels of pollution as proxy for an individual's exposure to pollution. This approach may be oversimplified because individuals can undertake avoidance activities to reduce the effect of pollution: households can respond to an increasing level of particulate, for instance, by avoiding exposure or mitigating the effects of the exposure once they occur (Cropper and Oates, 1992). If people respond to a higher fine particulate concentration by staying inside or by increasing the avoidance behavior or by mitigating the effects, for instance, through curative care to the point that health actually improves, not controlling for this aspect may yield estimates that are lower bounds of the true effect (Neidell, 2004). So, a limitation of our study is the absence of proxy for individual avoidance or mitigating behavior in the data set.

References

- [1] E. Bakhshi, et al.,(2008), " The Positive Association Between Number of Children and Obesity in Iranian Women and Men: Results from the National Health Survey", *BMC Public Health* 2008, **8**:213-219.
- [2] S. Balia, A.M.Jones, (2008), " Mortality, Lifestyle and Socio-Economic Status", *Journal of Health Economics*, **27**:1-26.
- [3] X.Basagaca, et al.(2004), "Socioeconomic Status and Asthma Prevalence in Young Adults: The European Community Respiratory Health Survey", *American Journal of Epidemiology*, 160, No 2.
- [4] Bascom et al.(1996) Health effects of outdoor air pollution: part I, *American Journal of Respiratory and Critical Care Medicine*, 153: 477-98.
- [5] E.D. Bateman et al.(2008), "Global strategy for asthma management and prevention: GINA executive summary", *European Respiratory*, 31:143-178.
- [6] N.B. Belloc , L.Breslow (1972), " Relationship of Physical Health Status and Health Practices", *Preventive Medicine*, **1**:409-421.
- [7] M.Berger, J.P.Leigh, (1989), "Schooling, Self-Selection, and Health", *Journal of Human Resources*, **24**: 433-455.
- [8] J.R. Blaylock, W.N. Blisard (1992), "Self-evaluated health status and smoking behaviour", *Applied Economics*, **24**: 429--435.
- [9] M. Boezen, et al. (1998) "Peak expiratory flow variability, bronchial responsiveness, and susceptibility to ambient air pollution in adults". *American Journal of Respiratory and Critical Care Medicine*, 158: 1848--1854.
- [10] K. Bolin, B. Lindgren (2002), " Asthma and Allergy: the Significance of Chronic Conditions for Individual Health Behavior", *Allergy*, **57**: 115-122.
- [11] S. Braman (2006), "The Global Burden of Asthma", *CHEST*, 130 :4S-12S
- [12] Brook, et al. (2004) "Air Pollution and Cardiovascular Disease: a Statement for Healthcare Professionals from the Expert Panel on Population and Prevention Science of the American Heart Association" *Circulation* **109**: 2655--2671.
- [13] B. Brunekreef, S.T. Holgate (2002), " Air Pollution and Health", *Lancet*, **360**: 1233--1242.
- [14] R.V. Burkhauser, J. Cawley , M.D. Schmeiser (2009), "The timing of the rise in U.S. obesity varies with measure of fatness", *Economics & Human Biology*, 7: 307--318.
- [15] L.Cappellari S. P Jenkins (2003), " Multivariate Probit Regression Using Simulated Maximum Likelihood". *The Stata Journal*, **3**:278-294.
- [16] Y.Chen et al. (1999). Increased Effects of Smoking and Obesity on Asthma among Female Canadians: The National Population Health Survey, 1994--1995. *American Journal of Epidemiology*, 150: 255--262.
- [17] P.Contoyannis, A. M Jones, (2004), "Socio-Economic Status, Health and Lifestyle", *Journal of Health Economics*, **23**:965-995
- [18] M.L. Cropper (1981), " Measuring the Benefits from Reduced Morbidity", *The American Economic Review*, **71**: 235-240.
- [19] M.L. Cropper, W.E. Oates (1992), " Environmental Economics: A Survey", *Journal of Economic Literature*, **30**: 675-740.
- [20] M.J. Daniels, et al. (2000), " Estimating particulate matter-mortality dose-response curves and threshold levels: an analysis of daily time-series for the 20 largest US

- cities", *American Journal of Epidemiology*, **152**: 397-406.
- [21] R. De Marco et al. (2000) "Differences in Incidence of Reported Asthma Related to Age in Men and Women", *American Journal of Respiratory Critical Care Medicine*, **162**: 68-74.
- [22] P.V. Dicpinigaitis, K. Rauf (1998), "The influence of gender on cough reflex sensitivity", *CHEST*, **13**:1319--132.
- [23] C. Di Novi (2010), "The Influence of Traffic-Related Pollution on Individuals' Lifestyle: Results from the BRFSS", *Health Economics*, **19**: 1318-1344.
- [24] R.K. Dishman, J. F. Sallis (1994), "Determinants and Interventions for Physical Activity and Exercise". In: *Physical activity, fitness and health: International proceedings and consensus statement*, C. Bouchard, R. J. Shephard, and T. Stephens (Eds.). Champaign, IL: Human Kinetics, pp. 214–238.
- [25] F. Dominici et al.(2002), "Air Pollution and Mortality: Estimating Regional and National Dose-response Relationships", *Journal American Stat Association*, **97**:100--111.
- [26] H.Dume, S.K.Weiland,U.Keil (1998), "Epidemiological Analysesof the Relationship between Environmental Pollution and Asthma", *Toxicology Letters*,102-103, 307-316.
- [27] T.M.L. Eagan, et al. (2002), " Incidence of asthma and respiratory symptoms by sex, age and smoking in a community study ", *European Respiratory Journal*, 19: 599-605.
- [28] P. Ernst (2002), "Environmental Measures and Asthma", *Chest*, 122:1509-1510
- [29] Erbsland M, Ried W, Ulrich V (1994), "Health, Health Care, and the Environment. Econometric Evidence from German Micro Data." *Health Economics* **4**:169-182.
- [30] D. Erzen et al. (1997) "Income Level and Asthma Prevalence and Care Patterns". *American Journal of Respiratory Critical Care Medicine* **155**: 1060-1065.
- [31] J. Espinosa , WN Evans, (2008), "Heightened mortality after the death of a spouse: marriage protection or marriage selection?", *Journal of Health Economics*,**27**:1326-1342...
- [32] E. S. Ford et al. (2003), "Self-Reported Asthma and Health- Related Quality of Life: Findings From the Behavioral Risk Factor Surveillance System", *CHEST*, 123:119-127.
- [33] Global Initiative for Asthma (GINA), National Heart, Lung and Blood Institute (NHLBI). Global strategy for asthma management and prevention. Global Initiative for Asthma (GINA), National Heart, Lung and Blood Institute (NHLBI): Bethesda (MD); 2006. p. 339. Available from: www.ginasthma.com.
- [34] W. W. Gould (1993) Linear splines and piecewise linear functions. *Stata Technical Bulletin* **15**:13–17.
- [35] G.A. Grassman (1996), "Obtaining Information about Susceptibility from Epidemiological Literature", *Toxicology*, 111: 253-270.
- [36] M. Grossman, (1972), " On the Concept of Health Capital and the Demand for Health", *Journal of Political Economy*, **80**: 223-255.
- [37] Y. Hu, B. Wolfe (2002),"Health Inequality between Black and White Women". Institute for Research on Poverty Discussion Paper no. 1251-02.
- [38] A.P. Jones (1998), "Asthma and Domestic Air Quality", *Social Science and Medicine*, **47**: 755-764.
- [39] Katsouyanni K, Touloumi G, Spix C. (1997), "Short-term Effects of Ambient Sulphur Dioxide and Particulate Matter on Mortality in 12 European Cities: Results from Times

- Series Data from the APHEA Project" *British Medical Journal*, **314**: 1658--63.
- [40] DS. Kenkel (1991), " The Demand for Preventive Medical Care", *Applied Economics*, **26**: 313-325.
- [41] A.C. King, G.Brassington, (1997), "Enhancing Physical and Psychological Functioning in Older Family Caregivers: the Role of Regular Physical Activity. *Annual Behavioral Medicine*, **19** :91-100
- [42] G.S. Maddala (1983), "Limited Dependent and Qualitative Variables in Econometrics", Cambridge University Press.
- [43] B.H. Marcus, et al., (1999), "The Efficacy of Exercise as an Aid for Smoking Cessation in Women: a Randomized Controlled Trial", *Archives of Internal Medicine* , **159**:1229-34.
- [44] Masoli, M, Fabian, D, Holt, S, et al. (2004) The global burden of asthma: executive summary of the GINA Dissemination Committee report. *Allergy* 59,469-478.
- [45] J.Mullahy, J. (1999): " It'll Only Hurt a Second: Microeconomic Determinants of Who Gets Flu Shots," *Health. Economics*, 8, 9-24.
- [46] M.Neidell (2004), "Air Pollution, Health, and Socio-Economic Status: the Effect of Outdoor Air Quality on Childhood Asthma". *Journal of Health Economics*, **23**:1209-1236.
- [47] Office of Management and Budget, "Standards for Defining Metropolitan and Micropolitan Statistical Areas; Notice".Federal Register, Vol. 65, No. 249, December 2000.
- [48] Physical Activity Guidelines for Americans. 2008. U.S. Department of Health and Human Services, www.health.gov/paguidelines.
- [49] S. Roberts (2004), "Biologically Plausible Particulate Air Pollution Mortality Concentration–Response Functions", *Environmental Health Perspectives*, 112: 309-313.
- [50] Samet JM, Zeger SL, Berhane K. Particulate air pollution and daily mortality: replication and validation of selected studies. The Phase I Report of the Particle Epidemiology Evaluation Project. Cambridge, MA:Health Effects Institute.
- [51] L.Sandvik, et al. (2003), " Physical Fitness as a Predictor of Mortality Among Healthy, Middle-Aged Norwegian Men", *New England Journal of Medicine*, **328**:533-537.
- [52] Schwartz, J., Weiss, S.T., (1994a). The relationship of dietary fish intake to level of pulmonary function in the first National Health and Nutrition Survey (NHANES I). *European.Respiratory Journal*, **7**:1821-1824.
- [53] Schwartz, J., Weiss, S.T., (1994b). Relationship between dietary vitamin C intake and pulmonary function in the First National Health and Nutrition Examination Survey (NHANES I). *American Journal of Clinical Nutrition*. 59, 110-114.
- [54] G.Schwarz (1978) " Estimating the Dimension of a Model", *Annals of Statistics* 6: 461–464.
- [55] K.Sexton (1997), "Sociodemographic Aspects of Human Susceptibility to Toxic Chemicals: Do Class and Race Matter for Realistic Risk Assessment?" *Environmental Toxicology and Pharmacology*, **4**:261-269.
- [56] E. N.Sherwood, R. W. J effery (2000) " The Behavioral Determinants of Exercise: Implications for Physical Activity Interventions" *Annual Review of Nutrition*, **20**: 21-44.
- [57] M. R. Stolley, M.L.Fitzgibbon (1997) Effects of an Obesity Prevention Program on the

- Eating Behavior of African American Mothers and Daughters, *Health Education & Behavior*, Vol. 24, No. 2, 152-164
- [58] A.Takeda et al. (2004) " Multigenerational Family Structure in Japanese Society: Impacts on Stress and Health Behaviors Among Women and Men", *Social Science and Medicine*, **59**:69-81.
- [59] United States Environmental Protection Agency, 2003, Criteria Pollutants-Metropolitan Area Trends, <http://www.epa.gov/oar/airtrends/metro.html>.
- [60] A. Valavanidis, T. Vlachogianni, and K. Fiotakis (2009), "Tobacco Smoke: Involvement of Reactive Oxygen Species and Stable Free Radicals in Mechanisms of Oxidative Damage, Carcinogenesis and Synergistic Effects with Other Respirable Particles", *International Journal Environmental Research and Public Health*; **6**: 445--462.
- [61] S. Walters et al., 1995 Hospital Admission Rates for Asthma and Respiratory Disease in the West Midlands: Their Relationship to Air Pollution Levels, *Thorax*, **50**: 948--54.
- [62] J.P.Watson et al., 1996, "The Relationship Between Asthma Admission Rates, Routes of Admission, and Socioeconomic Deprivation", *European Respiratory Journal*, **9**:2087-2093.
- [63] Wen XJ, Balluz L, Mokdad A. (2009), "Association between media alerts of air quality index and change of outdoor activity among adult asthma in six states, BRFSS, 2005. *Journal Community Health*; **34**:40-46.
- [64] WHO. 1999. Environmental Health Indicators: Framework and Methodologies. Geneva:World Health Organization. Available: http://whqlibdoc.who.int/hq/1999/WHO_SDE_OEH_99.10.pdf [accessed 16 March 2010].

Appendix I: Tables

TABLE I: Variable Name and Definition

<i>Variables Name</i>	<i>Variables Definition</i>
asthma	1 if has asthma, 0 otherwise
smoke	1 if has smoked at least 100 cigarettes in his life and is current smoker, 0 otherwise
alcohol	1 if is at risk for heavy drinking, 0 otherwise
diet	1 if consumes fruits/vegetables at least once per day, 0 otherwise
execise	1 if meets recommendations for physical activity, 0 otherwise
obese	1 if is at risk for overweight or obesity (BMI >25.0000), 0 otherwise
flushot	1 if has flu shot in the 12 months before the interview, 0 otherwise
stress	1 if mental health (including stress) was not good, 0 otherwise
inactive	1 if is inactive, 0 otherwise
male	1 if male 0 otherwise
age	Age in years
hmo	1 if has health care coverage , 0 otherwise
element	1 if elementary school or Kindergarden, 0 otherwise
high_sch	1 if attend high school or high school graduate, 0 otherwise
collg	1 if attend college or college graduate, 0 otherwise
married	1 if married, 0 otherwise
divorce	1 if divorced, 0 otherwise
widow	1 if widow, 0 otherwise
seprd	1 if sepatated,0 otherwise
never_married	1 if never married, 0 otherwise
unmar_couple	1 if member of an unmarried couple, 0 otherwise
out_labor_market	1 if retired, homemaker, student, 0 otherwise
unemployed	1 if unemployed, 0 otherwise
self_emp	1 if employed or self-employed, 0 otherwise
white	1 if White, 0 otherwise
black	1 if Black, 0 otherwise
other_race	1 if other race, 0 otherwise
nummen	Number of men living in household
numwomen	Number of women living in household
children	Number of children less than 18 years of age living in household
physhlth	1 if during the past 30 days physical health was not good, 0 otherwise
pm2.5_aqi	maximum daily PM2.5 AQI

TABLE II a): Summary Statistics

	Means	St. Deviation
asthma	0.1172	0.3216
age	46.6242	17.0868
male	0.4074	0.4914
smoke	0.2470	0.4313
alcohol	0.0518	0.2216
diet	0.9643	0.1854
exercise	0.2212	0.4150
obese	0.2231	0.4163
flu shot	0.3089	0.4621
stress	0.3596	0.4799
inactive	0.2521	0.4342
hmo	0.9018	0.2976
element	0.0271	0.1623
high_sch	0.375	0.4842
collg	0.3001	0.4584
married	0.5176	0.4997
divorce	0.1469	0.3540
widow	0.1	0.3
seprd	0.0222	0.1473
never_married	0.1843	0.3878
unmar_couple	0.0289	0.1678
employed	0.6368	0.4809
out labor market	0.3302	0.4703
unemployed	0.0329	0.1786
white	0.8361	0.3702
black	0.1071	0.3093
other_race	0.0568	0.2315

TABLE II b): Summary Statistics

Metropolitan Area Name	Maximum Daily PM 2.5 AQI
Chicago,IL	144
Cincinnati,OH-KY	132
Cleveland-Lorain	155
Columbus,OH	142
Des Moines,IA	92
Huntington-Ashla	134
Las Vegas,NV-AZ	132
Louisville,KY-IN	141
Memphis,TN-AR-MS	154
Milwaukee-Waukes	161
Minneapolis-St.	118
Nashville,TN	97
Omaha,NE-IA	122
Phoenix-Mesa,AZ	146
St, Louis,MO-IL	143
Tucson,AZ	62

TABLE III a) Estimated Coefficients of Asthma- PM_{2.5} Model with Threshold*

	asthma	smoke	alcohol	exercise	diet	obese	flu shot	stress
age	-0.0068 (0.000)	0.0085 (0.000)	-0.0037 (0.118)	-0.0150 (0.000)	0.0104 (0.000)	0.0059(0.000)	0.0264 (0.000)	-0.0208 (
male	-0.3484(0.000)							
physhlth	0.3364 (0.000)							
smoke	0.2346 (0.003)							
alcohol	0.4246 (0.017)							
exercise	-0.0641 (0.482)							
diet	0.0555 (0.761)							
obese	0.2834 (0.002)							
flu_shot	0.3999 (0.000)							
stress	0.1407 (0.088)							
inactive		0.1096 (0.003)	- 0.1105 (0.090)	-1.067 (0.000)	-0.4166 (0.000)	0.2605 (0.000)	-0.0641 (0.111)	0.1044 (
hmo	0.0019 (0.981)	-0.2174(0.000)	-0.03784 (0.658)	0.0254 (0.691)	0.1061 (0.232)	-0.1012 (0.086)	0.3558 (0.000)	-0.0348 (
high_school	-0.0585 (0.276)	0.1648 (0.000)	0.0221 (0.720)	-0.1252 (0.007)	-0.1563 (0.027)	0.0788 (0.054)	0.0229 (0.577)	-0.0654 (
collg	-0.0392 (0.514)	-0.3161(0.000)	-0.1953 (0.004)	0.1914 (0.000)	0.1859 (0.032)	-0.1417 (0.002)	0.1275 (0.003)	-0.1459 (
divorce		0.2490 (0.000)	0.3084(0.000)	0.0596 (0.309)	-0.2984 (0.001)	-0.0104 (0.849)	-0.0828 (0.131)	0.1930 (
widow		-0.1143 (0.074)	0.0145 (0.906)	-0.0758 (0.406)	-0.0115(0.939)	-0.1546(0.031)	-0.1403(0.036)	0.1521 (C
seprd		0.1801(0.096)	0.1396 (0.464)	0.0286 (0.828)	-0.0922 (0.644)	0.1726 (0.131)	-0.0498 (0.687)	0.2001 (C
never_married		-0.0437 (0.397)	0.2603 (0.001)	-0.0129(0.825)	-0.3928 (0.000)	-0.0552 (0.337)	0.1609 (0.005)	0.0271 (C
unmar_couple		0.3199 (0.001)	0.4395 (0.001)	-0.1525 (0.168)	-0.1784 (0.289)	-0.1045 (0.343)	-0.0480 (0.681)	0.2838 (C
out_labor_market	0.1616 (0.003)	-0.1067(0.006)	-0.2338 (0.001)	-0.1299(0.005)	0.0867 (0.267)	-0.1008 (0.021)	0.2042 (0.000)	0.1097 (C
unemployed	-0.0351(0.787)	0.0602 (0.503)	-0.0333 (0.813)	-0.2452 (0.028)	0.1356 (0.385)	0.1689 (0.070)	0.0277 (0.786)	0.3285 (C
black	0.0481(0.517)	-0.2812 (0.000)	-0.3342 (0.001)	0.0365 (0.564)	-0.0235 (0.799)	0.3055 (0.000)	0.0117 (0.840)	-0.1877 (
other_race	-0.0214 (0.832)	-0.2397 (0.001)	-0.1705 (0.135)	0.0895 (0.256)	0.0187 (0.877)	0.2560 (0.001)	-0.0697(0.382)	-0.0401 (
children		-0.0574(0.138)	-0.2039 (0.002)	-0.0809 (0.066)	0.0175 (0.812)	0.0863 (0.044)	-0.1407 (0.001)	-0.0255 (
nummen		0.1052 (0.000)	0.0816 (0.064)	0.1163 (0.000)	0.0896 (0.092)	-0.0357 (0.266)	-0.0302 (0.351)	-0.0925 (
numwomen		-0.0900 (0.004)	0.000(0.998)	-0.0829 (0.022)	0.1258 (0.026)	0.0077 (0.821)	0.0310(0.359)	0.0861 (C
pm2.5_aqi<100	0.0016 (0.571)	-0.0045 (0.017)	-0.0118 (0.000)	-0.0075 (0.001)	0.0117 (0.002)	0.0014 (0.522)	0.0052 (0.013)	-0.0081 (
pm2.5_aqi>100	0.0012 (0.389)	0.0029 (0.002)	0.0065 (0.000)	0.0035 (0.002)	-0.0068 (0.001)	-0.000 (0.834)	-0.0018 (0.074)	0.0034 (

* p-values in parentheses.

TABLE III b) Estimated Coefficients of Asthma- PM_{2.5} Linear Model without Threshold*

	asthma	smoke	alcohol	exercise	diet	obese	flu shot	stress
age	-0.0069 (0.000)	0.0087 (0.000)	-0.0033 (0.171)	-0.0148 (0.000)	0.0099 (0.000)	0.0059 (0.000)	0.0261 (0.000)	-0.0205 (0.000)
male	-0.3486 (0.000)							
physlth	0.3364 (0.000)							
smoke	0.2337 (0.003)							
alcohol	0.4129 (0.022)							
exercise	-0.0693 (0.446)							
diet	0.0618 (0.734)							
obese	0.2839 (0.002)							
flu_shot	0.4004 (0.000)							
stress	0.1345 (0.102)							
inactive		0.1029 (0.005)	-0.1235 (0.057)	-1.0736 (0.000)	0.3992 (0.000)	0.2619 (0.000)	0.0578 (0.150)	-0.0939 (0.013)
hmo	0.0018 (0.982)	-0.2213 (0.000)	-0.0472 (0.579)	0.0194 (0.762)	0.1103 (0.212)	-0.1004 (0.088)	0.3584 (0.000)	-0.0406 (0.457)
high_school	-0.0585 (0.276)	0.1611 (0.000)	0.0152 (0.805)	-0.1314 (0.004)	-0.1432 (0.042)	0.0794 (0.052)	0.0254 (0.537)	-0.0704 (0.068)
collg	-0.0399 (0.506)	-0.3109 (0.000)	-0.1843 (0.007)	0.1969 (0.000)	0.1809 (0.036)	-0.1431 (0.002)	0.1221 (0.005)	-0.1377 (0.001)
divorce		0.2526 (0.000)	0.3111 (0.000)	0.0649 (0.268)	-0.3072 (0.001)	-0.0112 (0.837)	-0.0869 (0.113)	0.1981 (0.000)
widow		-0.1132 (0.077)	0.0148 (0.904)	-0.0733 (0.421)	-0.0145 (0.923)	-0.1549 (0.030)	-0.1417 (0.034)	0.1536 (0.025)
seprd		0.1823 (0.092)	0.1326 (0.487)	0.0307 (0.815)	-0.0977 (0.623)	0.1718 (0.132)	-0.0522 (0.673)	0.2037 (0.063)
never_married		-0.0366 (0.478)	0.2765 (0.001)	-0.0033 (0.955)	-0.4084 (0.000)	-0.0568 (0.322)	0.1534 (0.008)	0.0377 (0.467)
unmar_couple		0.3374 (0.000)	0.4764 (0.000)	-0.1329 (0.230)	-0.2240 (0.180)	-0.1089 (0.322)	-0.0654 (0.576)	0.3098 (0.001)
out_labor_market	0.1612 (0.003)	-0.1029 (0.009)	-0.2272 (0.001)	-0.1225 (0.008)	0.0802 (0.302)	-0.1015 (0.020)	0.2009 (0.000)	0.1147 (0.004)
unemployed	-0.0352 (0.787)	0.0696 (0.438)	-0.0208 (0.883)	-0.2313 (0.039)	0.1233 (0.431)	0.1668 (0.073)	0.0178 (0.862)	0.3418 (0.000)
black	0.0467 (0.527)	-0.2693 (0.000)	-0.3043 (0.002)	0.0521 (0.409)	-0.0559 (0.539)	0.3032 (0.000)	0.0017 (0.977)	-0.1700 (0.002)
other_race	-0.0232(0.816)	-0.2087(0.002)	-0.1057 (0.349)	0.1334 (0.087)	-0.0599 (0.613)	0.2492 (0.001)	-0.0974 (0.218)	0.0090 (0.896)
children		-0.0580 (0.134)	-0.2036 (0.002)	-0.0831 (0.059)	0.0177 (0.808)	0.0864 (0.044)	-0.1401 (0.001)	-0.0269 (0.491)
nummen		0.1067 (0.000)	0.0839 (0.056)	0.1184 (0.000)	0.0820 (0.120)	-0.0361(0.260)	-0.0321(0.322)	-0.0897 (0.002)
numwomen		-0.0899 (0.004)	0.0041 (0.930)	-0.0816 (0.024)	0.1176 (0.036)	0.0077 (0.822)	0.0312 (0.356)	0.0864 (0.006)
pm2.5_aqi	0.0013 (0.095)	0.0006 (0.212)	0.0007 (0.424)	0.0001(0.835)	-0.0012 (0.298)	0.0003 (0.679)	0.0003 (0.671)	-0.0000 (0.930)

* p-values in parentheses.

TABLE IV a): Estimated Correlation Coefficients of Asthma- PM_{2.5} Model with Threshold*

	asthma	smoke	alcohol	exercise	diet	obese	flu shot	stress
asthma	1							
smoke	-0.0448 (0.269)	1						
alcohol	-0.1242 (0.076)	0.2672 (0.000)	1					
exercise	0.0269 (0.536)	-0.0199 (0.365)	0.0247 (0.391)	1				
diet	-0.0611(0.372)	-0.0940 (0.012)	-0.0792 (0.121)	0.0530 (0.202)	1			
obese	-0.0259 (0.063)	-0.0301 (0.144)	-0.0189 (0.501)	-0.1197 (0.000)	-0.0118 (0.695)	1		
flu shot	-0.0725 (0.121)	-0.0089 (0.658)	-0.0421 (0.125)	-0.0361 (0.112)	0.0821(0.006)	0.0198 (0.363)	1	
stress	-0.0181 (0.660)	0.0653 (0.001)	0.0811 (0.002)	-0.0156 (0.463)	-0.0396 (0.147)	0.0806 (0.000)	-0.0267 (0.191)	1

TABLE IV b): Estimated Correlation Coefficients of Asthma- PM_{2.5} Linear Model without Threshold*

	asthma	smoke	alcohol	exercise	diet	obese	flu shot	stress
asthma	1							
smoke	-0.0439(0.279)	1						
alcohol	-0.1181 (0.095)	0.2673(0.000)	1					
exercise	0.0298 (0.492)	-0.0172 (0.435)	0.0279 (0.333)	1				
diet	-0.0638 (0.349)	-0.0970 (0.009)	-0.0851 (0.095)	0.0509 (0.220)	1			
obese	-0.0262 (0.059)	-0.0306 (0.137)	-0.0196 (0.486)	-0.1201 (0.000)	-0.0123 (0.682)	1		
flu shot	-0.0727 (0.119)	-0.0106 (0.601)	-0.0445 (0.103)	-0.0383 (0.092)	0.0832 (0.005)	0.0202 (0.356)	1	
stress	-0.0143 (0.728)	0.0677 (0.000)	0.0848 (0.001)	-0.0129 (0.546)	-0.0418 (0.125)	0.0791 (0.000)	-0.0293 (0.151)	1

* p-values in parentheses.




Figure 1. Relation between pollution and investments in health.

1. Solimene L., *Market Failures and State Intervention*
2. Solimene L., *The Efficiency under Private and Public Ownership: Some Empirical Evidence*
3. Baici E., Dell'Aringa C., *The EMS Effect on the Italian Labour Market*
4. Lucifora C., *Union Density and Relative Wages: Is there a Relationship?*
5. Lucifora C., Sestito P., *Determinazione del salario in Italia: una rassegna della letteratura empirica*
6. Martini G., *Testing Different Bargaining Theories: A Pilot Experiment*
7. Lucifora C., Rappelli F., *Profili retributivi e carriere: un'analisi su dati longitudinali*
8. Dell'Aringa C., Lucifora C., *Wage Dispersion and Unionism: Are Unions Egalitarian?*
9. Martini G., *Horizontal Price Fixing and Antitrust Policy: A Sequentially Rational Design*
10. Cassuti G., Dell'Aringa C., Lucifora C., *Labour Turnover and Unionism*
11. Solimene L., *Regolamentazione ed incentivi all'innovazione nel settore delle telecomunicazioni*
12. Bigard A., Guillotin Y., Lucifora C. e F. Rappelli, *An International Comparison of Earnings Mobility: The Case of Italy and France*
13. Martini G., *Laboratory Tests of a Kinked Demand Curve Model with Discounting and Game-theoretic Foundations*
14. Martini G., *A Multi-period Antitrust Game: The Dynamic Effects of Competition Policy*
15. Piccirilli G., *Monetary Business Cycles with Imperfect Competition and Endogenous Growth*
16. Dell'Aringa C., *Pay Determination in the Public Service: An International Comparison*
17. Lucifora C., *Rules Versus Bargaining: Pay Determination in the Italian Public Sector*
18. Piccirilli G., *Hours and Employment in a Stochastic Model of the Firm*
19. Cappellari L., *The Covariance Structure of Italian Male Wages, 1974 – 1988*
20. Lucifora C., *Working Pooors? An Analysis of Low Wage Employment in Italy*
21. Lucifora C., Origo F., *Alla ricerca della flessibilità: un'analisi della curva dei salari in Italia*
22. Dell'Aringa C., Vignocchi C., *Employment and Wage Determination for Municipal Workers: The Italian Case*
23. Cappellari L., *Wage Inequality Dynamics in the Italian Labour Market: Permanent Changes or Transitory Fluctuations?*
24. Cappellari L., *Low-pay transitions and attrition bias in Italy: a simulated maximum likelihood approach*
25. Pontarollo E., Vitali F., *La gestione del parco tecnologico elettromedicale tra outsourcing e integrazione verticale*
26. Cappellari L., *Do the 'Working Pooors' Stay Poor? An Analysis of Low-Pay Dynamics in Italy*
27. Dell'Aringa C., Lucifora C., *Inside the black box: labour market institutions, wage formation and unemployment in Italy*
28. Filippini L., Martini G., *Vertical Differentiation and Innovation Adoption*
29. Lucifora C., Simmons R., *Superstar Effects in Italian Football: an Empirical Analysis*
30. Brunello G., Lucifora C., Winter-Ebmer R., *The Wage Expectations of European College Students*

31. Cappellari L., *Earnings dynamic and uncertainty in Italy: How do they differ between the private and public sectors?*
32. Piccirilli G., *Unions and Workforce Adjustment Costs*
33. Dell'Aringa C., *The Italian Labour Market: Problems and Prospects*
34. Bryson A., Cappellari L., Lucifora C., *Does Union Membership Really Reduce Job Satisfaction?*
35. Cappellari L., *The effects of high school choices on academic performance and early labour market outcomes*
36. Cappellari L., Jenkins S. P., *Transitions between unemployment and low pay*
37. Dell'Aringa C., Pagani L., *Collective Bargaining and Wage Dispersion*
38. Comi S., *University enrolment, family income and gender in Italy*
39. Ghinetti P., *The Wage Effect of Working in the Public Sector When Education and Sector Choices Are Endogenous: An Empirical Investigation for Italy*
40. Piccirilli G., *Unions, Job Protection and Employment*
41. Bryson A., Cappellari L., Lucifora C., *Why so unhappy? The effects of unionisation on job satisfaction*
42. Brunello G., Cappellari L., *The Labour Market Effects of Alma Mater: Evidence from Italy*
43. Dell'Aringa C., Pagani L., *Regional Wage Differentials and Collective Bargaining in Italy*
44. Dell'Aringa C., *Industrial Relations and Macroeconomic Performance*
45. Prandini A., *Structural Separation or Integration in Italian Fixed Tlc: Regulatory and Competition Issues*
46. Ghinetti P., *The Public-Private Job Satisfaction Differential in Italy*
47. Cappellari L., Ghinetti P., Turati G., *On Time and Money Donations*
48. Cappellari L., Leonardi M., *Earnings Instability and Tenure*
49. Cappellari L., Dorsett R., Haile G., *State dependence, duration dependence and unobserved heterogeneity in the employment transitions of the over-50s*
50. Piccirilli G., *Job protection, industrial relations and employment*
51. Cappellari L., Lucifora C., *The "Bologna Process" and College Enrolment Decisions*
52. Piccirilli G., *Contingent Worksharing*
53. Ursino G., *Supply Chain Control: A Theory of Vertical Integration*
54. Barron G., Ursino G., *Underweighting Rare Events in Experience Based Decisions: Beyond Sample Error*
55. Comi S., *Family influence on early career outcomes in seven European countries*
56. Cottini E., Lucifora C., *Health and Low-pay: a European Perspective*
57. Comi S., *Intergenerational mobility in seven European Countries*
58. Dell'Aringa C., Pagani L., *Labour Market Assimilation and Over Education: The Case of Immigrant Workers in Italy*
59. Cappellari L., Tatsiramos K., *Friends' Networks and Job finding Rates*
60. Cappellari L., Dell'Aringa C., Leonardi M., *Flexible Employment Job Flows and Labour Productivity*
61. Fedele A., Tedeschi P., *Quality and Reputation: Is Competition Beneficial to Consumers?*
62. Di Novi C., *The Indirect Effect of Fine Particulate Matter on Health through Individuals' Life-style*